		, ,1	eren part of the 4 A DEC
FORM!	PTO-139 1-2000)	90 (Modified) U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER
	TI	RANSMITTAL LETTER TO THE UNITED STATES	PU-9925
		DESIGNATED/ELECTED OFFICE (DO/EO/US)	U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR
		CONCERNING A FILING UNDER 35 U.S.C. 371	To be 1s 1 n/d 0 1 8 0 4 2
NTEI		IONAL APPLICATION NO. INTERNATIONAL FILING DATE PCT/SE00/01174 June 21, 2000	PRIORITY DATE CLAIMED June 30, 1999
Com	bina	NVENTION torial Library with Particle Classes that can be Distinguished by Tw	vo Features e.g. Size, Density, Color, etc.
		T(S) FOR DO/EO/US "man	
Appli	cant l	herewith submits to the United States Designated/Elected Office (DO/EO/US) the	ne following items and other information:
1.	\boxtimes	This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.	
2.		This is a SECOND or SUBSEQUENT submission of items concerning a filin	1
3.	\boxtimes	This is an express request to begin national examination procedures (35 U.S.C	
4.	\boxtimes	(9) and (24) indicated below.The US has been elected by the expiration of 19 months from the priority date	(Article 31).
5.	\boxtimes	A copy of the International Application as filed (35 U.S.C. 371 (c) (2))	ζ
٥.		a. ⊠ is attached hereto (required only if not communicated by the Interna	tional Bureau)
		b. \square has been communicated by the International Bureau.	
		c. \square is not required, as the application was filed in the United States Rece	viving Office (RO/US)
6.		An English language translation of the International Application as filed (35 U	- '
0.		a. \square is attached hereto.	7.3.C. 371(C)(2)).
		b. has been previously submitted under 35 U.S.C. 154(d)(4).	
7.		Amendments to the claims of the International Application under PCT Article	19 (35 H S C 371 (c)(3))
,.		a. are attached hereto (required only if not communicated by the International Approximation and International A	
		b. \square have been communicated by the International Bureau.	anonai Burcau).
		c. \square have not been made; however, the time limit for making such amenda	ments has NOT expired
		d. \square have not been made and will not be made.	ments has to t expired.
8.		An English language translation of the amendments to the claims under PCT A	Article 19 (35 U.S.C. 371(c)(3))
9.	\boxtimes	An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).	There 17 (33 0.3.C. 371(c)(3)).
10.		An English language translation of the annexes to the International Preliminary Article 36 (35 U.S.C. 371 (c)(5)).	y Examination Report under PCT
11.	\boxtimes	A copy of the International Preliminary Examination Report (PCT/IPEA/409).	}
12.	\boxtimes	A copy of the International Search Report (PCT/ISA/210).	
It	ems 1	13 to 20 below concern document(s) or information included:	
13.	\boxtimes	An Information Disclosure Statement under 37 CFR 1.97 and 1.98.	
14.		An assignment document for recording. A separate cover sheet in compliance	with 37 CFR 3.28 and 3.31 is included.
15.	\boxtimes	A FIRST preliminary amendment.	
16.		A SECOND or SUBSEQUENT preliminary amendment.	ì
17.		A substitute specification.	1
18.		A change of power of attorney and/or address letter.	
19.		A computer-readable form of the sequence listing in accordance with PCT Rul	le 13ter.2 and 35 U.S.C. 1.821 - 1.825.
20.		A second copy of the published international application under 35 U.S.C. 154((d)(4).
21.		A second copy of the English language translation of the international applicat	ion under 35 U.S.C. 154(d)(4).
22.	\boxtimes	Certificate of Mailing by Express Mail	1
23.	\boxtimes	Other items or information:	
		copy of this transmittal letter for charging purposes return postcard	
		·	

J607 Res'd PCT/PTO 1 0 DEC 2001

U.S. APPLICATION	NO (IF KNOWN	ATTORNEY'S DOCKET NUMBER										
To	pedsyig dd]		PU-9925									
724. The fol	lowing fees are	submitt	ted:.					CAL	CULATIONS	PTO USE ONLY		
BASIC NATIONA												
international	mational prelimits search fee (37 cional Search Rej	\$1040.00										
☐ International USPTO but	l preliminary ex International Se	\$890.00										
but internati	l preliminary ext onal search fee (\$740.00										
but all claim	l preliminary exa s did not satisfy	\$710.00										
International and all claim	l preliminary ex- ns satisfied provi	amınatı isions c	on fee (37) of PCT Art	CFR 1.482) paid to U icle 33(1)-(4)	SPIO		\$100.00					
	ENTER	APPI	ROPRI	ATE BASIC FE	E AMO	OUN	NT =		\$1,040.00			
Surcharge of \$130.0 months from the ear	liest claimed pr	iority o	date (37 CI	R 1.492 (e)).	☐ 20		□ 30		\$0.00			
CLAIMS	NUMB	ER FII	LED	NUMBER EXT	RA		RATE					
Total claims	16		- 20 =	0		X	\$18.00		\$0.00			
Independent claims			- 3 =	0		Х	\$84.00		\$0.00			
Multiple Dependent	Claims (check			ADOVE CALC	TIT AT	LON.			\$0.00	· · · · · · · · · · · · · · · · · · ·		
	11			ABOVE CALC			<u> </u>		\$1,040.00			
Applicant clair reduced by 1/2		status. S	See 37 CFI	R 1.27). The fees indic					\$0.00			
					SUB	ГОТ	$\Gamma AL =$		\$1,040.00			
Processing fee of \$1 months from the ear	130.00 for furnis liest claimed pr	shing th	ne English date (37 CI	translation later than FR 1.492 (f)).	□ 20	0	□ 30 +		\$0.00			
2		· · · · · · · · · · · · · · · · · · ·		TOTAL NAT	IONAI	LFE	EE =		\$1,040.00			
Fee for recording the accompanied by an	e enclosed assig	nment er sheet	(37 CFR 1 t (37 CFR	.21(h)). The assignme 3.28, 3.31) (check if a	ent must b	e).			\$0.00			
				TOTAL FEES	ENCL	OSI	E D =		\$1,040.00			
									int to be: efunded	\$		
								_		\$		
a. \square A ch	neck in the amou	ınt of		to cover the	above fee	s is er	nclosed.	-				
	se charge my Do			o500-588	in the am			40.00	to cover the	e above fees.		
	Commissioner i		y authoriz 500-58 8	ed to charge any additi A duplicate cop				quired,	or credit any ov	verpayment		
d. 🗌 Fees	are to be charg	ed to a	credit card	. WARNING: Inform on this form. Provide	ation on t	this fo	rm may beco					
NOTE: Where an	appropriate tir	ne limi	t under 37	CFR 1.494 or 1.495	has not b	een n				J-2036.		
		_	d to restor	e the application to p	ending st	atus.		2		2 1		
SEND ALL CORRI	ESPONDENCE	TO:			Home Man							
Royal N. Ronning Amersham Biosci	ences			:	SIGNAZURE SIGNAZURE							
800 Centennial Av Piscataway, New						Royal N. Ronning, Jr.						
i iscataway, ivew	dersey noods					NA	ME					
(732) 457-8423						32	,529					
						RE	GISTRATIC	ON NU	MBER	,		
						De	cember 10	, 2001				
1						$\frac{D}{DA}$,				
						21						

10/018042 JEO7 Rec'd PCT/PTO 10 DEC 2001

PU-9925



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application of:

N. Norrman

Group Art Unit:

To be assigned

Serial Number:

To be assigned

Examiner:

To be assigned

Filing Date:

December 7, 2001

Title:

Combinatorial Library with Particle Classes that can be

Distinguished by Two Features e.g. Size, Density, Color, etc.

FIRST PRELIMINARY AMENDMENT

Honorable Assistant Commissioner of Patents Box Patent Application Washington, D.C. 20231

Sir:

Please consider the following amendments and remarks in connection with the prosecution of the captioned application, which is a filing under 35 U.S.C. § 371 and claims priority to international application number PCT/SE00/01174 filed June 21, 2000. This application also claims priority to patent application number 9902479-6 filed in Sweden on June 30, 1999.

In the Claims

Please amend claim 1 as follows:

1. (once amended) A method of identifying one or more substances having affinity for a given target substance, comprising:

(i) providing a set of particle classes, each said particle class being distinguishable from [another class] other classes within the set by at least one physical property, each particle class being comprised of at least two [subclasses] sub-classes, wherein each [subclass] sub-class is distinguishable from [another subclass] the other sub-classes of said particle class by [another] a physical property, which is different from the property which distinguishes the particle class from other classes within the set, and further wherein [different from the property of said particle classes,]each [subclass comprising] sub-class contains particles having at least one of said one or more substances attached to the surface thereof as a ligand, said ligands being different from ligands attached to particles of other particle classes or [subclasses] sub-classes;

(iii) combining a plurality of <u>classes or sub-classes</u> to form at least one mixture,

(iii) distributing [the mixtures] <u>said at least one mixture</u> in separate vessels;

(iv) exposing [each] <u>said mixture in said separate vessels</u> to said target substance;

(v) [washing away] <u>removing</u> all target substance not having bound to any ligand;

and

(iv) identifying to which particle [sub-class(es)]classes or sub-classes said target substance actually has (have) bound, by[the following steps:] identifying in which vessel or vessels target substance has bound to particles in the mixture present in said vessel or vessels, identifying to which particle class said target substance has bound; and

identifying to which particle <u>class or</u> sub-class said target substance has bound.

Please amend claim 2 as follows:

2. (once amended) The method [as claimed in]of claim 1, wherein each particle class is characterized by one of the <u>physical</u> properties [in]selected from the group consisting of size, density, color and shape.

Please amend claim 3 as follows:

3. (once amended) The method [as claimed in]of claim 2, wherein each particle subclass is characterized by one of the <u>physical</u> properties [in the group]selected from the group consisting of size, density, color and shape, but is difference from the property characterizing the particle class.

Please amend claim 4 as follows:

4. (once amended) The method [as claimed in]of claim 1, wherein the mixtures are formed by mixing two particle classes or sub-classes in a ratio such that the difference in [number of one class with respect to the other is detectable in the identifying step]the relative amount of each class or sub-class can be used to determine the particle class to which the target substance is bound.

Please amend claim 5 as follows:

5. (once amended) The method [as claimed in any preceding claim]of claim 1, wherein the target substance is marked so as to be detectable.

Please amend claim 6 as follows:

6. (once amended) The method [as claimed in]of claim 5, wherein the [marking is performed]target substance is marked by attaching a moiety selected from the group consisting of a fluorescent moiety, a radioactive moiety, a colored moiety.

Please amend claim 7 as follows:

7. (once amended) The method [as claimed in]of claim 5, wherein the target substance reacts with the ligand to which it binds to provide a detectable effect, such as fluorescence or color.

Please amend claim 8 as follows:

8. (once amended) The method [as claimed in any preceding claim]of claim 1, wherein the identification is performed by ocular inspection under microscope.

Please amend claim 9 as follows:

9. (once amended) The method [as claimed any preceding claim]of claim 1, wherein the identification is performed by image analysis with a computer.

Please amend claim10 as follows:

10. (once amended) The method [as claimed in any preceding claim]of claim 1, wherein said mixture is exposed to at least two target substances.

Please amend claim 11 as follows:

11. (once amended) A library of different ligands, comprising particles belonging to a plurality of <u>particle</u> classes, each particle class being distinguishable from [another class]the other particle classes by at least one physically distinguishable property, each particle class being comprised of at least two [subclasses]subclasses, wherein each [subclass]sub-class is distinguishable from [another subclass]the other sub-classes by [another]a physical property of the particle class from the property which distinguishes the particle class from other classes within the set, and further wherein[, different from the property(ies) of said particle classes,] those particles belonging to [one and]the same sub-class having at lease one type of ligand attached to their surface.

Please amend claim 12 as follows:

12. (once amended) The ligand library [as claimed in]of claim 11, wherein one of said properties of the particle classes or sub-classes is the size[, suitably the diameter] of the particles.

Please amend claim 13 as follows:

13. (once amended) The ligand library [as claimed in]of claim 11, wherein one of said properties of the particle classes or sub-classes is the density of the particle.

Please amend claim 14 as follows:

14. (once amended) The ligand library [as claimed in]of claim 11, wherein one of said properties of the particle classes or sub-classes is the shape of the particle.

Please amend claim 15 as follows:

15. (once amended) The ligand library [as claimed in]of claim 11, wherein one of said properties of the particle classes or sub-classes is the color of the particle.

Remarks

Claims 1-16 are pending in the instant application. Applicants have amended claims 1-15 to more fully conform with U.S. practice and to delete multiple dependencies. A version of the claims marked up to show the amendments, as well as a clean version of the claims encompassing the amendments, is attached hereto.

Applicants respectfully assert that all amendments are fairly based on the specification, and respectfully request their entry.

Applicants believe that the claims, as amended, are in allowable form, and earnestly solicit the allowance of claims 1-16.

Respectfully submitted,

Royal N. Ronning, Jr. 32,529

Attorney for Applicants

Amersham Biosciences 800 Centennial Avenue P. O. Box 1327 Piscataway, New Jersey 08855-1327

Tel: (732) 457-8423

Fax: (732) 457-8463

Claims (marked-up version showing amendment(s))

- 1. (once amended) A method of identifying one or more substances having affinity for a given target substance, comprising:
 - (i) providing a set of particle classes, each said particle class being distinguishable from [another class]other classes within the set by at least one physical property, each particle class being comprised of at least two [subclasses]sub-classes, wherein each [subclass]sub-class is distinguishable from [another subclass]the other sub-classes of said particle class by [another]a physical property, which is different from the property which distinguishes the particle class from other classes within the set, and further wherein [different from the property of said particle classes,]each [subclass comprising]sub-class contains particles having at least one of said one or more substances attached to the surface thereof as a ligand, said ligands being different from ligands attached to particles of other particle classes or [subclasses]sub-classes;
 - (ii) combining a plurality of <u>classes or</u> sub-classes to form at least one mixture,

 (iii) distributing [the mixtures]said at least one mixture in separate vessels;

 (iv) exposing [each]said mixture in said separate vessels to said target substance;

 (v) [washing away]removing all target substance not having bound to any ligand;
 - (v) [washing away]removing all target substance not having bound to any ligand; and
 - (iv) identifying to which particle [sub-class(es)]classes or sub-classes said target substance actually has (have) bound, by[the following steps:] identifying in which vessel or vessels target substance has bound to particles

in the mixture present in said vessel or vessels, identifying to which particle class said target substance has bound; and identifying to which particle <u>class or sub-class said target substance</u> has bound.

- 2. (once amended) The method [as claimed in]of claim 1, wherein each particle class is characterized by one of the <u>physical</u> properties [in]selected from the group consisting of size, density, color and shape.
- 3. (once amended) The method [as claimed in]of claim 2, wherein each particle subclass is characterized by one of the <u>physical</u> properties [in the group]selected from the group consisting of size, density, color and shape, but is difference from the property characterizing the particle class.
- 4. (once amended) The method [as claimed in]of claim 1, wherein the mixtures are formed by mixing two particle classes or sub-classes in a ratio such that the difference in [number of one class with respect to the other is detectable in the identifying step]the relative amount of each class or sub-class can be used to determine the particle class to which the target substance is bound.
- 5. (once amended) The method [as claimed in any preceding claim]of claim 1, wherein the target substance is marked so as to be detectable.

- 6. (once amended) The method [as claimed in]of claim 5, wherein the [marking is performed]target substance is marked by attaching a moiety selected from the group consisting of a fluorescent moiety, a radioactive moiety, a colored moiety.
- 7. (once amended) The method [as claimed in]of claim 5, wherein the target substance reacts with the ligand to which it binds to provide a detectable effect, such as fluorescence or color.
- 8. (once amended) The method [as claimed in any preceding claim] of claim 1, wherein the identification is performed by ocular inspection under microscope.
- 9. (once amended) The method [as claimed any preceding claim] of claim 1, wherein the identification is performed by image analysis with a computer.
- 10. (once amended) The method [as claimed in any preceding claim]of claim 1, wherein said mixture is exposed to at least two target substances.
- 11. (once amended) A library of different ligands, comprising particles belonging to a plurality of <u>particle</u> classes, each particle class being distinguishable from [another class]the other particle classes by at least one physically distinguishable property, each particle class being comprised of at least two [subclasses]subclasses, wherein each [subclass]subclass is distinguishable from [another subclass]the other subclasses by [another]a physical property of the particle class

from the property which distinguishes the particle class from other classes within the set, and further wherein[, different from the property(ies) of said particle classes,] those particles belonging to [one and]the same sub-class having at lease one type of ligand attached to their surface.

- 12. (once amended) The ligand library [as claimed in]of claim 11, wherein one of said properties of the particle classes or sub-classes is the size[, suitably the diameter] of the particles.
- 13. (once amended) The ligand library [as claimed in]of claim 11, wherein one of said properties of the particle classes or sub-classes is the density of the particle.
- 14. (once amended) The ligand library [as claimed in]of claim 11, wherein one of said properties of the particle classes or sub-classes is the shape of the particle.
- 15. (once amended) The ligand library [as claimed in]of claim 11, wherein one of said properties of the particle classes or sub-classes is the color of the particle.

Claims (clean version encompassing amendments)

- 1. (once amended) A method of identifying one or more substances having affinity for a given target substance, comprising:
 - (i) providing a set of particle classes, each said particle class being distinguishable from other classes within the set by at least one physical property, each particle class being comprised of at least two sub-classes, wherein each sub-class is distinguishable from the other sub-classes of said particle class by a physical property, which is different from the property which distinguishes the particle class from other classes within the set, and further wherein each sub-class contains particles having at least one of said one or more substances attached to the surface thereof as a ligand, said ligands being different from ligands attached to particles of other particle classes or sub-classes;
 - (ii) combining a plurality of classes or sub-classes to form at least one mixture,
 - (iii) distributing said at least one mixture in separate vessels;
 - (iv) exposing said mixture in said separate vessels to said target substance;
 - (v) removing all target substance not having bound to any ligand; and
 - (iv) identifying to which particle classes or sub-classes said target substance actually has (have) bound, by

identifying in which vessel or vessels target substance has bound to particles in the mixture present in said vessel or vessels, identifying to which particle class said target substance has bound; and

identifying to which particle class or sub-class said target substance has bound.

- (once amended) The method of claim 1, wherein each particle class is characterized by one of the physical properties selected from the group consisting of size, density, color and shape.
- 3. (once amended) The method of claim 2, wherein each particle sub-class is characterized by one of the physical properties selected from the group consisting of size, density, color and shape, but is difference from the property characterizing the particle class.
- 4. (once amended) The method of claim 1, wherein the mixtures are formed by mixing two particle classes or sub-classes in a ratio such that the difference in the relative amount of each class or sub-class can be used to determine the particle class to which the target substance is bound.
- 5. (once amended) The method of claim 1, wherein the target substance is marked so as to be detectable.
- 6. (once amended) The method of claim 5, wherein the target substance is marked by attaching a moiety selected from the group consisting of a fluorescent moiety, a radioactive moiety, a colored moiety.

- 7. (once amended) The method of claim 5, wherein the target substance reacts with the ligand to which it binds to provide a detectable effect, such as fluorescence or color.
- 8. (once amended) The method of claim 1, wherein the identification is performed by ocular inspection under microscope.
- 9. (once amended) The method of claim 1, wherein the identification is performed by image analysis with a computer.
- 10. (once amended) The method of claim 1, wherein said mixture is exposed to at least two target substances.
- 11. (once amended) A library of different ligands, comprising particles belonging to a plurality of particle classes, each particle class being distinguishable from the other particle classes by at least one physically distinguishable property, each particle class being comprised of at least two sub-classes, wherein each sub-class is distinguishable from the other sub-classes by a physical property of the particle class from the property which distinguishes the particle class from other classes within the set, and further wherein those particles belonging to the same sub-class having at lease one type of ligand attached to their surface.

- 12. (once amended) The ligand library of claim 11, wherein one of said properties of the particle classes or sub-classes is the size of the particles.
- 13. (once amended) The ligand library of claim 11, wherein one of said properties of the particle classes or sub-classes is the density of the particle.
- 14. (once amended) The ligand library of claim 11, wherein one of said properties of the particle classes or sub-classes is the shape of the particle.
- 15. (once amended) The ligand library of claim 11, wherein one of said properties of the particle classes or sub-classes is the color of the particle.
- 16. The use of a ligand library as claimed in claim 11 for screening purposes.

10

15

20

25

30

wo 01/01141 4/PRTS

10/018042 JC07 Rec'd PCT/PTO 1 0 DEC 2001 PCT/SE00/01174

1

COMBINATORIAL LIBRARY WITH PARTICLE CLASSES THAT CAN BE DISTINGUISHED BY TWO FEATURES E.G. SIZE, DENSITY, COLOR, ETC.

The present invention relates generally to screening methods and selective identification of target substances.

Background of the invention

In the pharmaceutical area i.a. it is often necessary to test very large numbers of substances against one or more target compounds to find out if any of the tested substances possess affinity towards the target, or in some other way interacts with the target. This normally is performed by binding each candidate compound (as a ligand) to some matrix e.g. a particle, loading a number of the particles in one reaction vessel of some kind, and incubating with the target substance. Either the candidate compound (ligand) or the target is marked with identifiable groups such as fluorescent or radioactive groups or nuclei. After the reaction is deemed to have come to completion analysis of the vessel is performed to see if the marker substance or group is present, indicating that the target has indeed reacted with the substance bound to the matrix.

Screening large numbers of substances of course requires large numbers of vessels, which is laborious and tedious to handle. 10000 compounds would require 10000 individual reaction vessels to be handled.

Another possibility is to mark particles with say 10000 different ligands. A mixture is prepared comprising particles having 100 different ligands and the mixture is placed in a well of a microtiter plate. 100 such mixtures are prepared totaling 10000 different ligands and said mixtures are placed in one well each. All wells are exposed to and incubated with the target substance marked with a fluorescent moiety. If fluorescence is detected in one well one knows that a hit is present in that well. A second experiment is performed where the 100 different ligands in said well are individually placed in one well each, and the incubation and exposure is repeated. The well exhibiting fluorescence will then contain the desired substance. Thus, it is necessary to perform two sets of experiments.

Therefore, it would be desirable to have access to methods and means for rendering such screening procedures less time and space consuming, and to enable the procedure to be carried out in one step.

- 5 US-5,858,670 relates to identification by sequencing of peptides bound to beads. The method requires many operations. It mentions staining of beads (col. 41, lines 47-54) for identification. This particular aspect per se, does not form a part of the present invention
- In "One-Bead-One-Structure Combinatorial Libraries", <u>Biopolymers</u>, Vol. 37, 177-198, there is disclosed a very complicated (many steps required) method of making libraries for the identification of binding to specific ligands. The actual screening procedure requires a large number of iterations.
- US-5,817,751 discloses libraries where beads have "identifier tags". These tags can be
 "microscopically or otherwise distinguishable features", (size, shape, mass, charge, color).
 See col. 23, lines 10-21, and lines 42-46. The screening process involves releasing the tags from the beads and subsequent amplification thereof.
- EP 0 773 227 A1 discloses "identifier tags", meaning some "physical attribute" (examples given are size, shape, color, optical density) by which a solid support (e.g. a bead) can be identified and distinguished. See p 5, lines 17-33. The screening procedure is conventional, and involves sorting out fluorescing beads by using a cell sorting instrument (FACS).
- US-5,162,863 discloses the possibility of using particles of different sizes for identification purposes.

Summary of the Invention

The present invention seeks to provide a method of identifying one or more substances having affinity for a given target substance, that would require a smaller number of reaction vessels than what is necessary today, and also a smaller number of operations, thereby speeding up the process of investigation substantially.

This object is achieved by the method according to the invention, whereby a combination of distinguishable properties of particles to which ligands are attached, is used for identification of those particles to which a target substance has bound. The method is defined in claim 1.

- By providing a class of particles having one property in common, and subdividing each such class into sub-classes having another property, it becomes possible to screen a very large number in only two steps.
- In a second aspect of the invention there is provided a library of ligands, comprising
 individually distinguishable particles of a matrix material, having ligands attached to the
 surface, and being suitable for binding candidate compounds (ligands), and usable in
 screening procedures. This aspect is defined in claim 11.
- A third aspect of the invention is the use in screening procedures, of a ligand library

 comprising individually distinguishable particles of a matrix material, suitable for binding

 candidate compounds, which is characterized by using fact that the particles are

 distinguishable, as a marker for each said candidate compound. The library is defined in claim

 16.
- Furthermore, in preferred embodiments, selecting the distinguishable property to be different particle size in combination with density, is very powerful in this respect.

Brief Description of the Drawings

- 25 The invention will now be described with reference to examples and to the drawings, in which
 - Fig. 1 is a schematic representation of a micro titer well in which a mixture of 100 different particle-ligand combinations has been incubated;
- Fig. 2 is another schematic representation of a micro titer well;
 - Fig. 3 is photograph of the contents of a microtiter well taken through a fluorescence microscope; and

Fig. 4 is a graphical output from an image analysis.

Detailed Description of Preferred Embodiments

10

5

For the purpose of this application, the term "particle class" is taken to mean particles having at least one property distinguishing them from other particles of another "class". One such property can be size (diameter), another the density. Still another kind of "particle class" is formed when a number of particles of one class having a first ligand attached, is used together with particles of the same class but having a second ligand attached thereto, but where the ratio of the number of particles having different ligands is different, e.g. 2:1.

A further particle class is formed when a coordinate is assigned to a set of particles

20

The term "sub-class" is taken to mean particles having as common features, the features or 15 properties of a "particle class", and having been treated in some way so as to be distinguishable from another "sub-class". A "sub-class" is e.g. formed when particles of one size are provided with one ligand, and another "sub-class" is formed when the same particles are provided with a second ligand, different from the first.

Within one and the same class, i.e. particles having the same size, "sub-classes" are also formed by providing particles with e.g. different density but having the same diameter.

25

30

The term "ligand" shall mean one moiety attached to a particle, or a group of such moieties.

The basic idea behind the invention is the insight that in a mixture of objects, wherein groups of objects in said mixture have some distinguishable and identifiable feature in common within each group, it is easy to identify each group of objects. Also it is easy to identify if there has been a change in the property of the objects in one group. If for example a mixture consists of all white golf balls, tennis balls and footballs, it is very easy by visual inspection to determine that say the golf balls have changed color to green.

10

15

20

25

30

This insight has been applied by the inventor in the field of screening for substances having some desired property amongst large numbers of candidate substances, or ligands.

An extremely simplified example is the case where two ligands are to be examined for affinity to a target substance suitably marked, e.g. with a fluorescent marker. Each ligand would then be bound to particles with mutually different size and then the particles are mixed. The mixture is placed in well of a micro titer plate and incubated. After incubation the well is washed. If the target selectively binds to one ligand it will be an easy matter to identify which ligand has bound since one can measure the size of the fluorescing particles and thereby determine which the ligand is.

As another simplified and illustrative but slightly more complex example, let us assume that one is interested in testing four ligands for affinity to some macro-molecule, e.g. a protein. Let us also assume that each different ligand is individually bound to a particle. Furthermore we assume that the particles are of only two different classes, in this case we assume two different particle sizes. Two mixtures are prepared, each comprising two sets of particles having distinguishable size, and having different ligands bound thereto. The two mixtures are placed in one well each and incubated with the macro-molecule, appropriately marked with e.g. a fluorescent marker. Thus, we have two wells each containing the same two different particle classes, but having mutually different ligands bound to them. After incubation the wells are washed to remove non-bound substances. If the macro-molecule binds to one ligand it will be a straight forward matter to identify which one by determining which particle size and which well exhibits fluorescence. This is most easily done under a fluorescence microscope, and can be done by ocular inspection in a case like the described, where there are only few wells and few ligands present. Other methods of detection are available, e.g. so called flow cytometry in combination with Coulter-counter techniques. These methods are well known to the skilled man and need not be discussed in detail herein.

For production screening using large numbers of ligands and particles in large numbers of wells, image analysis by computer using specialized software is more feasible. An example of such commercial software is LEICA® Q 500 MC.

20

25

30

The skilled man will appreciate that the principle demonstrated above is applicable to larger numbers of different particle sizes and large numbers of wells, as will be explained by further examples below.

The basic principle of a first embodiment of the invention may be said to encompass identification of a ligand having reacted with a target substance, by two parameters, namely a) particle size, and b) coordinate for the particle on the micro titer plate (simply in which well it is present).

Still another property of the particles that is usable as an identifying "marker" is the density. It is relatively easy to make particles having well defined densities. If ligands are bound to such particles, mixtures of said particles can be separated by centrifugation in a density gradient, and fluorescent bands in the test tube after centrifugation can be easily determined, and the position along the tube will indicate which density fraction is a hit.

The basic principle can be further elaborated by making mixtures containing different number ratios between particles of the same size having different ligands, and providing them in the same well. Thus, if two different ligands are bound to one and the same particle size but in two fractions, and these two fractions are mixed e.g. in a ratio 1:2, of course it will be an easy

matter to determine to which fraction the target has bound by the fluorescence intensity being either 1/3 or 2/3 of the maximum possible, if all particles would have fluoresced.

The particles to be used may be made of any material to which suitable ligands of interest can be attached or bound. The particles are preferably hydrophilic and built up of one or more polymers which are insoluble in water. Hydrophobic polymers that have been derivatized to become hydrophilic are included. Suitable polymers are polyhydroxy polymers, e.g. based on polysaccharides, such as agarose, dextran, cellulose, starch, pullulan, etc. and completely synthetic polymers, such as polyacrylic amide, polymethacrylic amide, poly(hydroxyalkylvinyl ethers), poly(hydroxyalkylacrylates) and polymethacrylates (e.g. polyglycidylmethacrylate), polyvinylalcohols and polymers based on styrenes and divinylbenzenes, and copolymers in which two or more of the monomers corresponding to the above-mentioned polymers are included. Polymers, which are soluble in water, may be derivatized to become insoluble, e.g. by cross-linking and by coupling to an insoluble body

via adsorption or covalent binding. Hydrophilic groups can be introduced on hydrophobic polymers (e.g. on copolymers of monovinyl and divinylbenzene) by polymerization of monomers exhibiting groups which can be converted to OH, or by hydrophilization of the final polymer, e.g. by adsorption of suitable compounds, such as hydrophilic polymers.

5

The particles can also be based on inorganic material, such as silica. Preferred particles lack hydrolytically unstable groups, such as silane, ester and amide groups.

The particles may be porous.

10

The term "hydrophilic particle" in practice means that the accessible surface of the particles is hydrophilic in the sense that can be penetrated by aqueous liquids. Typically the accessible surfaces on a hydrophilic particle expose a plurality of polar groups for instance comprising oxygen and/or nitrogen atoms. Examples of such polar groups are hydroxy, amino, carboxy, ester, ether of lower alkyls (such as (-CH₂CH₂O-)_nH where n is an integer).

15

Suitable size fractions of such particles are obtained by sieving, using standard methods well known to the skilled man. The resulting particle fractions will have some spread in terms of average diameter, but the spread can be controlled so that overlap between fractions can be adequately controlled.

20

Furthermore, a certain overlap in fraction size is no problem since the image analysis software that calculates average particle sizes is able to distinguish with a high degree of certainty to which fraction the fluorescing particles belong.

25

Another possible material for the particles is poly-styrene. By using this material it is possible to make mono-disperse particles without overlap in fraction size.

30

In order to prepare suitable mixtures when the number of particles and ligands is large, so called factor design is preferably used. The theory behind this methodology is disclosed in "Statistics for Experiments" by Box et al, ISBN 0-471-09315-7.

The invention will now be further described by way of the following non-limiting examples.

EXAMPLES

Example 1

5

10

15

20

25

30

10 particle fractions ranging in size from fraction 1 of 10-20 μm up to fraction 10 of 100-110 μm are prepared. Each size fraction is further subdivided into 10 density fractions ranging from 1,01 up to 1,19 with increments of 0,02. Thus, in all 100 different sub classes of particles defined by both a) size and b) density is produced. To particles of each sub class a different ligand is bound to provide 100 unique combinations of ligand and particle.

10000 different ligands are to be tested. Thus, the 100 particle subclasses are used to provide 100 sets of 100 particle-ligand combinations. Each set of 100 combinations is placed in one well each of a 100-well microtiter plate. Then all wells are incubated with a target compound. The compound is suitably marked with a fluorescent moiety. In addition to being identifiable by the two properties of the particle (size and density), each ligand is identified by a coordinate, i.e. the well number.

After incubation the wells are washed to remove all target compounds that have not bound to any ligands.

Fig. 1 shows a matrix symbolizing one well in which it has been determined that the size fraction 70-80 µm exhibits fluorescence (symbolized by the particles being filled), and thus contains a subclass of particles to which the target compound has bound. Of course only 10% of the particles of this size will fluoresce, but it cannot be decide which ligand has reacted.

In order to determine which density fraction contains the target bound to the particle, the entire contents in the well is centrifuged in a density gradient, so as to yield bands corresponding to each density. The band (density 1,11) containing the fraction having target bound thereto will then fluoresce. This is symbolized by the dots representing the density fraction in question being filled.

The cross section of the two rows will identify the particular ligand that has bound to the target molecule.

Example 2

5

In order to double the number of possible ligands to test (or alternatively to reduce the number of wells needed for the screening), it is also possible to use the number ratio between subclasses of particles as a marker.

Thus, 10 particle fractions ranging in size from fraction 1 of 10-20 μm up to fraction 10 of 100-110 μm are prepared. Each size fraction is further subdivided into 10 density fractions ranging from 1,01 up to 1,19 with increments of 0,02. Thus, in all 100 different sub classes of particles defined by both a) size and b) density is produced. To particles of each sub class a different ligand is bound to provide 100 unique combinations of ligand and particle.

15

20000 different ligands are to be tested. To this end, the 100 particle subclasses are used to provide a first lot of 100 sets of 100 particle-ligand combinations, which yields 10000 combinations. Furthermore, a second lot of 100 sets of 100 particle-ligand combinations, but with 10000 other ligands than in the first lot are made. In one and the same well one set of 100 particle-ligand combinations from the first lot is combined with twice as much (total number of particles or weight of the mixture or some other measure of quantity) from the second lot. This means that for each individual particle (having a unique size and density), there will be two possible ligands. The well is then incubated with a target molecule suitably marked.

25

30

20

After washing the wells to remove the non-bound target molecules, it is found that the size fraction 30-40 μ m exhibits fluorescence (see Fig 2 which is a schematic representation of the contents of the well). It is also determined by simple counting or by measuring a total intensity of the fluorescence, that the number of fluorescing particles is twice as many as would have been the case if the target had reacted with particles from the first lot, and thus it must be from the second lot.

A separation by density through centrifugation is performed as in Example 1, yielding 10 bands, one of which is fluorescing (density 1,01). Again, all information needed to determine which ligand has reacted is available.

5 Example 3

In this example it is demonstrated that the number of possible ligands to test can be increased also by mixing the particle classes in an intelligent way.

Particles of three different particle sizes, average diameters being 15 μm, 25 μm and 55 μm respectively are used and 18 different ligands (designated A-S) are bound to these particles, thus forming 54 different combinations of ligand/particle. Thus, a combination of ligand A with a particle having the diameter 15 μm is designated A-15, ligand B bound to a 55 μm particle is designated B-55 etc.

15

Mixtures of these ligand-particle aggregates are prepared according to the following protocol:

	Mixture No.	Composition (ligand/particle size)
	1	A-15; B-25; C-55; D-15; E-25; F-55
20	2	G-25; H-55; I-15; K-25; L-55; M-15
	3	N-55; O-15; P-25; Q-55; R-15; S-25
	4	A-15; B-25; C-55; D-25; E-55; F-15
	5	G-25; H-55; I-15; K-55; L-15; M-25
	6	N-55; O-15; P-25; Q-15; R-25; S-55

According to this protocol, a combination can occur either in one well only, or in two different wells. For example the combination A-15 occurs in well No. 1 and 4, whereas combination D-15 occurs only in well No. 1.

These mixtures are placed in one well each (designated with the same number as the mixture numbers) of a micro titer plate, thus in six wells. The mixtures are incubated with one target substance, suitably marked with a fluorescent moiety. After completed incubation the wells are washed with water to remove all unbound target substances.

It is found that wells No. 2 and 5 exhibited fluorescence, and that the fluorescent particles have a diameter of 25 μ m by image analysis. From this information it can be concluded that ligand G has bound to the target molecule, since the only combination with a particle of diameter 25 μ m common to well 2 and 5 is G-25. This being an extremely simple example, it is appreciated that the protocol of mixing can be much more sophisticated and involved, in order to distinguish one reacting species among a large number of possibilities.

In Fig. 3 a typical image viewed in a fluorescence microscope is shown. As can be clearly seen it is possible even with ocular inspection to distinguish three different particle sizes (in this case for illustrative purpose, all particles fluoresce in order to be able to see them; in a real run of course only one particle size should be visible). An image analysis by computer yields the result shown in Fig. 4, wherein also the distribution of sizes within each nominal class can be seen.

In a further development of the method, the mixtures of ligand-particle combinations are incubated with two or more target molecules. To this end the targets are suitably marked with distinguishable markers, such as fluorescent moieties exhibiting fluorescence of different wave lengths.

What is claimed is:

1. A method of identifying one or more substances having affinity for a given target substance, comprising:

5

providing a set of particle classes, each said particle class being distinguishable from another class by at least one physical property, each particle class being comprised of at least two subclasses, wherein each subclass is distinguishable from another subclass by another physical property, different from the property of said particle classes, each subclass comprising particles having at least one of said substances attached to the surface thereof as a ligand, said ligands being different from ligands attached to particles of other particle classes or subclasses;

combining a plurality of sub-classes to form at least one mixture,

15

10

distributing the mixtures in separate vessels;

exposing each mixture to said target substance;

20

washing away all target substance not having bound to any ligand; and

identifying to which particle sub-class(es) said target substance actually has(have) bound, by the following steps:

25

identifying in which vessel or vessels target substance has bound to particles

in the mixture present in said vessel or vessels, identifying to which particle class said target substance has bound; and

30

identifying to which particle sub-class said target substance has bound.

2. The method as claimed in claim 1, wherein each particle class is characterized by one of the properties in the group size, density, color and shape.

15

25

- 3. The method as claimed in claim 2, wherein each particle sub-class is characterized by one of the properties in the group size, density, color and shape, but is different from the property characterizing the particle class.
- 4. The method as claimed in claim 1, wherein the mixtures are formed by mixing two particle classes in a ratio such that the difference in number of one class with respect to the other is detectable in the identifying step.
- The method as claimed in any preceding claim, wherein the target substance is marked so as to be detectable.
 - 6. The method as claimed in claim 5, wherein the marking is performed by attaching a moiety selected from the group consisting of a fluorescent moiety, a radioactive moiety, a colored moiety.
 - 7. The method as claimed in claim 5, wherein the target substance reacts with the ligand to which it binds to provide a detectable effect, such as fluorescence or color.
- 20 8. The method as claimed in any preceding claim, wherein the identification is performed by ocular inspection under microscope.
 - 9. The method as claimed any preceding claim, wherein the identification is performed by image analysis with a computer.
 - 10. The method as claimed in any preceding claim, wherein said mixture is exposed to at least two target substances.
- 11. A library of different ligands, comprising particles belonging to a plurality of classes, each particle class being distinguishable from another class by at least one physically distinguishable property, each particle class being comprised of at least two subclasses, wherein each subclass is distinguishable from another subclass by another physical property,

14

different from the property(ies) of said particle classes, those particles belonging to one and the same sub-class having at least one type of ligand attached to their surface.

- 12. The ligand library as claimed in claim 11, wherein one of said properties of the particle classes or sub-classes is the size, suitably the diameter of the particles.
 - 13. The ligand library as claimed in claim 11, wherein one of said properties of the particle classes or sub-classes is the density of the particle.
- 10 14. The ligand library as claimed in claim 11, wherein one of said properties of the particle classes or sub-classes is the shape of the particle.
 - 15. The ligand library as claimed in claim 11, wherein one of said properties of the particle classes or sub-classes is the color of the particle.
 - 16. The use of a ligand library as claimed in claim 11 for screening purposes.

(19) World Intellectual Property Organization International Bureau





(43) International Publication Date 4 January 2001 (04.01.2001)

PCT

(10) International Publication Number WO 01/01141 A1

(51) International Patent Classification7: C12Q 1/68

(74) Agent: DR LUDWIG BRANN PATENTBYRÅ AB; Box 1344, S-751 43 Uppsala (SE).

(21) International Application Number:

PCT/SE00/01174

G01N 33/543,

(81) Designated States (national): JP, US.

(22) International Filing Date:

21 June 2000 (21.06.2000)

(84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

(25) Filing Language:

English

(26) Publication Language:

English

Published:

(30) Priority Data: 9902479-6

30 June 1999 (30.06.1999)

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

(71) Applicant (for all designated States except US): AMER-SHAM PHARMACIA BIOTECH AB [SE/SE]; Björkgatan 30, S-751 84 Uppsala (SE).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(72) Inventor; and

(75) Inventor/Applicant (for US only): NORRMAN, Nils [SE/SE]; Sunnerstavägen 26 j, S-756 51 Uppsala (SE).

(54) Title: COMBINATORIAL LIBRARY WITH PARTICLE CLASSES THAT CAN BE DISTINGUISHED BY TWO FEA-TURES E.G. SIZE, DENSITY, COLOR, ETC.

Particle size

		10-20		20-30		30-40		40-50		50-60		60-70		70-80		80	-90	90	-100	100	0-110
	1,01	0	а	0	þ	0	C	0	q	0	е	b	f		g	\Box) h	И) 1	\cup	
D	1,03	۰	k	0	1	0	B	0	ŋ	0	0	O	p		q	\cup	r	Γ) s	U	1
е	1,05	٥	ņ	٥	٧	0	х	0	У	0	z	0	a1 (•	a2	\Box	a3	C) a4	O	a 5
n	1,07	٥	a6	ò	a7	0	a 8	0	a9	0	b1	0	b2	5	b3	\subset) b4	\subset) b5	U	b6
s	1,09	0	b 7	0	b 8	0	b9	0	c1	0	c2	0	c3	5	c4	\mathbf{C}) c5) c6	\bigcirc	c 7
i	1,11	•	с8	•	с9	•	d١	•	d2	•	d3		d4 (d5		d6		d7		d8
t	1,13	0	d9	0	e1	0	e2	0	e3	0	e4	0	e5 (e6	\Box) e7	C) e8	\circ	e9
У	1,15		11	٥	12	0	fЗ	0	f4	0	f5	0	f6 f		f7	\Box) f8	C) f9	\bigcirc	g1
	1,17		g2	0	g3	0	g4	0	g5	0	g6	0	g7 !		g8	\subseteq) g9	\mathbf{C}) h1	\cup	h2
	1,19	٥	h3	0	h4	0	h5	0	h6	0	h7	0	h8		h9	\cup) 11	\Box) ;2	\bigcirc	13

(57) Abstract: The invention relates to a method of identifying one or more substances having affinity for a given target, comprising providing a set of particle classes, each said particle class being distinguishable from another class by at least one property. The particles belonging to one and the same class may have another property that distinguishes them, e.g. different densities, thus forming sub-classes. To each unique sub-class a number of unique ligands can be attached to sets of particles forming further sub-classes. A plurality of sub-classes is combined to form at least one mixture. The mixtures are distributed in separate vessels and exposed to said substance. All target substances not having bound to any candidate substance are washed away, and it is determined to which particle sub-class(es) said target substance actually has (have) bound.

a5 c7 c7 c9 e9 e9 h2 90-100 80-90 2 8 70-80 a1 b2 c3 c3 d4 e5 f6 f6 h8 **b**1 2828587 Particle size E × 88 61 59 E 45 30-40 c9 8 a/ 20-30 1 8 8 b 1 a 10-20 1,05 1,07 1,09 O e c s -- + >

Fig.

2/4






Fig. 2



F1g. 3

Size as a marker



Please type a plus sign (+) inside this box \longrightarrow +													
Under the Paperwork Reduced a valid OMB control number	Approved for use through 9/30/00 OMB 0651-0032 Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.												
			ket Number	PU-9925									
DECLARATION I		- 1	First Named Inventor Norrman										
PATENT AP	PLICATION		COMPLETE I	F KNOWN									
(37 CFI	R 1.63)	Application N	lumber 1	0 / 018	3,042								
		Filing Date	10-	10-Dec-2001									
☐ Declaration ☐ Submitted OR	Declaration Submitted after Initi	al Group Art Un	it To	To be assigned									
with Initial Filing	Filing (surcharge (37 CFR 1.16 (e)) required)	Examiner Na		be assigi									
As a below named invento	or, I hereby declare that:	**************************************											
My residence, post office ad	ddress, and citizenship are a	s stated below next to n	ny name.										
I believe I am the original, fin	rst and sole inventor (if only	one name is listed below	w) or an onginal, fi	st and joint invento	r (if plural								
	the subject matter which is clearly with Particle C												
Features e.g. Size	e, Density, Color, e	ic.	20 2.0 m. ga.	0.10 0 b y 1110									
the specification of which is attached hereto	(Title	of the Invention)											
OR x was filed on (MM/DD	MYYY) 06/21/2000) as Ur	nited States Applic	ation Number or Po	CT International								
Application Number PCT7	(C) (V) (G) (G) (G)	s amended on (MM/DE			(if applicable),								
I hereby state that I have revi	ewed and understand the co	ntents of the above ide	/	n, including the clai									
amended by any amendment I acknowledge the duty to dis	t specifically referred to abou	e.											
r dollowineage are duty to als	CIOSE INTOTTIBUIOTI WHICH IS III	aterial to paternaulity a	s delined in 37 CF	n 1.50.									
I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.													
Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed		y Attached? NO								
9902479-6	Sweden	06/30/1999											
1													
Additional foreign application	on numbers are listed on a s	upplemental pnonty dat	a sheet PTO/SB/0	2B attached hereto	:								
I hereby claim the benefit und Application Number(s			al application(s) list	ed below									
	, Fining Date	(MM/DD/YYYY)	numb suppl	ional provisional iers are listed on emental priority o SB/02B attached	a data sheet								

[Page 1 of 2]
Burden Hour Statement. This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time, you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED. FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

PTO/SB/01 (12-97) Please type a plus sign (+) inside this box + Approved for use through 9/30/00. OMB 0651-0032 Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number
--

DECLARATION — Utility or Design Patent Application

					3								
I hereby claim the benefit under 35 U.S.C. 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disc losed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose information which is maternal to patentability as defined in 37 CFR 1.56 which became available between the filing date of the and the national or PCT international filing date of this application.													
U.S. Par	ent Application or Number	PCT Paren		nt Filin N/DD/Y	ig Date YYY)		Parent Patent Number (if applicable)						
PCT/SE00/0	1174		06/2	1/2000	0		·						
As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application Ill Ill													
and Trademark Office connected therewith: Customer Number 22840													
OR Registered practitioner(s) name/registration number listed being An Label here													
Nam	ne		tration nber			Newy	ENT & TR A	UT(Regis	stration mber			
Additional registered	d practitioner(s) named or	n supplemental	Registered	Practition	er Inform	ation sheel	PTO/SB/	02C at	tached hereto	1.			
Direct all corresponde		ner Number Code Label	2284	40		OR	Cor	respon	ndence addr	ess below			
Name		·····											
Address										<u>- '</u>			
Address													
City				State	е		ZIP						
Country		Telephor	Telephone										
believed to be true; and	I statements made heren d further that these state apprisonment, or both, un t issued thereon.	ments were ma	ide with the	e knowled	ige that	willful false	statemer	nts and	the like so	o made are			
Name of Sole or I	First Inventor:			Пар	etition h	nas been f	iled for th	his uns	signed inven	ıtor			
Given Nar	me (first and middle (if	anyl)		Family Name or Surname									
Nils		, /		Norrman									
Inventor's Signature	Mil	Me	m						Date	14/1 -02			
Residence: City		State		Cou	ntry	SE			Citizenship	SE			
Post Office Address	Sunnersta	vagen 2	26 j										
Post Office Address	Uppsala,	Swede	n S-	756	51	SEX							
City	State		ZIP Country										
Additional inventor	ro oro bolog named an							0.00.0					